

VFC - Spoke wheel injury EN

spaakverwonding

Spoke-wheel injury

This kind of injury occurs when lower limbs get stuck in between spokes, frequently those of a bicycle. We did not see a fracture on the X-ray and there is no deep wound. A spoke-wheel injury is treated with a pressure bandage.

Summary

- Your child has a spoke-wheel injury of the foot.
- We did not see a fracture on the X-ray and there is no deep wound.
- A routine follow-up is not necessary for this type of injury.
- After 3 days, you can remove the band-aid and the pressure bandage.
- Thereafter, clean the wound every day by rinsing it in the shower until it has healed.
- You can find instructions on how to reapply the pressure bandage in the app.
- If you have any questions or problems during the recovery, do not hesitate to contact us on workdays from 09:00 – 11:00 AM – 2:00 – 4:00 PM via the Fracture Hot Line.

0-3 days

- Keep the band-aid on the wound.
- If the pressure bandage is loose, replace it with a tubigrip.
- Move your child's foot, when your child's pain allows this.

After 3 days

- Take care of your child's wound, follow the instructions listed underneath.
- Remove the pressure bandage and move your child's foot if your
- child's pain allows this.

Treatment

A spoke-wheel injury is common in the Netherlands. Your child will be treated with a band-aid and a pressure bandage.

Instructions

Follow these instructions for a good recovery.

- After 3 days, you can remove the band-aid and the pressure bandage.
- Your child can walk as soon as your child's pain allows it.
- When the pressure bandage is loose, you can replace this with a tubigrip. A tubigrip looks like a sock. Most likely, you will get one during your visit at the emergency department. You can also buy one at the pharmacy. Your child should not wear the tubigrip at night.
- At night, you can place a pillow underneath the foot to elevate it and thereby reduce swelling.
- Is your child in pain? Use a mild pain killer such as paracetamol (acetaminophen). Do not wait with the use of paracetamol until your child's pain is too much.

Instructions for the wound after 3 days

- Clean the wound every day, preferably in the shower. Continue cleaning the wound until it is closed. Put a new band-aid on the wound if necessary. Is the wound red and does the redness increase? Please contact the Fracture Hot Line. If a check-up seems necessary, we will schedule an appointment.

Recovery

- This type of injury almost always heals well with time and use – no specific treatment is required and routine follow-up is not necessary.
- Does the pain persist, or does the wound not heal? Please contact the Fracture Hot Line. If a check-up seems necessary, we will schedule an appointment.

Reapply the pressure bandage

In the video below you can find instructions on how to reapply the pressure bandage. You can buy new materials at a pharmacy or reuse the materials from the emergency department.

https://www.youtube.com/watch?v=bG-nGFrio_Q

Exercises

It is important that your child moves the foot and ankle as soon as the pain allows this. Specific exercises are not necessary.

Any questions?

Do you have any questions after reading this information? Ask your question via Laurentius hospital Fracture Hot Line.

Virtual Fracture Care App

Find more information about your injury in our app. The app also includes exercises that promote your recovery. You can find the app in the App Store (iPhone) or Google Store (Android).

You can also scan the QR code. You will then automatically be redirected to the app.



Contact

Fracture Hot Line

T 0475-38 22 11
(business days from 09:00 – 11:00 AM – 2:00 – 4:00 PM)

Source brochure: *Proefschrift Virtual Fracture Care: Direct ontslag vanaf de SEH voor patiënten met simpele stabiele letsels*

Laurentius Ziekenhuis
Mgr. Driessenstraat 6
(hoofdingang bereikbaar via de Oranjelaan)
6043 CV Roermond
www.lzr.nl

Patiëntencommunicatie Laurentius Ziekenhuis
VFC04 - Spoedeisende Hulp - oktober 2024